

Prevalence of Polycystic Ovary Syndrome (PCOS) In Relation to Body Mass Index (BMI) Variations: A Systematic Review

Syed Zaigham Ali Shah^{1*}, Muhammad Zubair², Muhammad Bilal³ and Mansoor Khan⁴

1,3,4MS Diagnostic Ultrasound, Ibadat International University Islamabad, Pakistan

2Lecturer, Department of Radiological Sciences & Medical Imaging, Ibadat International University Islamabad, Pakistan

Abstract

Background: Polycystic ovary syndrome (PCOS) is a serious clinical issue of the present age. It mostly affects women of reproductive age. Obesity is considered a potential risk factor associated with PCOS.

Objective: This systematic review was conducted to evaluate the prevalence of polycystic ovary syndrome (PCOS) in relation to body mass index (BMI) variations across different populations.

Methodology: A comprehensive search was performed using Google Scholar, Springer, and PubMed to identify relevant articles. The search was limited to studies published within the last 20 years (2005 - 2025). Only Studies reporting mean BMI, mean age, and prevalence of PCOS in the study population were chosen, while others were excluded.

Conclusion: BMI plays a crucial role in PCOS development and severity, increasing risks of metabolic and hormonal complications, emphasizing the need for standardized diagnosis and personalized treatment

Limitations: Access limitations and the scarcity of relevant studies posed challenges to this review.

Keywords: BMI, Infertility, Insulin resistance, Obesity, PCOS

Introduction

Polycystic ovary syndrome (PCOS) is a hormonal disorder marked by imbalance of sex hormones that primarily affects women of reproductive age [1,2]. Signs and symptoms that are associated with PCOS include irregular menstrual cycle, polycystic ovaries, hirsutism, obesity, acne, hyperandrogenism, insulin resistance, impaired glucose tolerance (Type 2 Diabetes) and infertility [2,3]. The majority of women with PCOS (38-88%) are either overweight or obese [4,5]. The level of obesity can be determined by using body mass index (BMI). It is calculated by dividing the weight in kilograms by the square of the height in meters. [6]. Adult BMI is classified into three categories: underweight ($<18.5 \text{ kg/m}^2$), normal weight (18.5 kg/m^2 to 24.9 kg/m^2), overweight (25 kg/m^2 to 29.9 kg/m^2), and obese ($\geq 30 \text{ kg/m}^2$) [7].

The etiology of PCOS has yet to be discovered and there is no exact known cause at the moment. However, a genetic component as well as several additional risk factors, such as obesity and insulin resistance, has been identified. [8]. In the current literature, there is a significant disparity in the BMI ranges associated with PCOS. Only a few researchers have looked into the relationship between BMI and PCOS. Due to variations with the diagnostic criteria for PCOS, these studies are frequently hampered by small sample sizes, selection bias, and are not comparable with the findings of other studies. The goal of this review is to evaluate the prevalence of PCOS based on ranges of BMI.

Literature Review

PCOS is a major clinical issue of the present age that has been reported to affect 8%-13% of reproductive aged women [9]. Obesity was first discovered to be a prevalent symptom of PCOS by Stein and Leventhal (1935) and later validated by many authors (Roger and Mitchell; 1952, Franks, 1989). Obesity has been estimated to affect up to 80% of PCOS women in the United States. [10]. According to a

recent Spanish study, PCOS is five times more common in overweight or obese women of childbearing age (28.3 percent) as compared to general population (5.5 percent) [11]. PCOS can appear at any time during a woman's reproductive life; however, it is most common during adolescence [12]. The incidence rate of infertility in reproductive-age women begins to climb at a BMI of 24 kg/m^2 and continues to increase as BMI rises [13]. According to a study published in June 2009, women with PCOS have reduced levels of follicle stimulating hormone (FSH) and increased levels of BMI as compared to normal women [14].

Association Between Obesity and PCOS

The most frequent finding in women with PCOS is obesity, and it is estimated that about 40%–80% of women suffering from this condition are either overweight or obese [15]. A recent systematic review of 35 studies found that overweight (BMI $25\text{--}30 \text{ kg/m}^2$) and obesity (BMI $\geq 30 \text{ kg/m}^2$) were about 2- and 2.8-fold more common in women with PCOS, respectively [16]. Obesity is linked to changes in adipokine and inflammatory cytokine production, which may contribute to obesity-related insulin resistance and metabolic syndrome [17]. In culture, hyper-insulinemia has been demonstrated to promote ovarian androgen synthesis by acting as a co-gonadotropin [18]. Patients with PCOS who have a BMI of 23 kg/m^2 or more are more likely to develop metabolic disorders [19]. With increasing weight, the prevalence of dyslipidemia rises [20]. A similar association persists in the metabolic syndrome [21]. However, a major multicenter research of women with PCOS reported no metabolic syndrome in those with PCOS and a BMI of 27 kg/m^2 [21].

It's unclear whether obesity causes PCOS or PCOS causes obesity. However, androgen exposure in postmenopausal women is the main cause of increasing visceral adiposity in obese as well as normal-weight women [22]. Similarly, a recent study suggests that uncontrolled obesity before puberty combined with severe insulin resis-

*Corresponding Author: *Syed Zaigham Ali Shah, Diagnostic Ultrasound, Ibadat International University Islamabad, Pakistan

Citation: Syed Zaigham Ali Shah*, Prevalence of Polycystic Ovary Syndrome (PCOS) In Relation to Body Mass Index (BMI) Variations: A Systematic Review. Jour of Clin & Med Case Rep, Imag 2026; 6(4): 1233.

tance can lead to the development of PCOS later in life (Figure 1) [23]. Obesity is also associated with elevated testosterone concentrations, suggesting increased androgen production [24]. Aromatase is a protein found in adipose tissue that promotes the synthesis of bio-active estrogens from androgens, which are then released into the bloodstream [25]. This can also cause elevated estrone levels, which have been observed in PCOS women [26]. According to a study published in 2011, elevated BMI levels are less specific than LAP (lipid accumulation product) in predicting IGT (impaired glucose tolerance) [27], confirming previous findings that LAP is better than BMI in predicting Type 2 Diabetes Mellitus. [28,29]. This could be explained by the fact that an increase in BMI can be induced by an increase in lean mass or an enlargement of the protective SAT (subcutaneous adipose tissue) in the lower extremities [30].




Figure 1: Schematic representation of development PCOS in women (Abbott et al., 2002).




Figure 2: Showing Role of Obesity in PCOS [31].

Material and Methods

This systematic review follows the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analyses) guidelines. A systematic review was conducted to evaluate variations in body mass index (BMI) among individuals diagnosed with PCOS across different populations. A comprehensive search was performed using Google Scholar, Springer, and PubMed to identify relevant articles. The search was limited to studies published within the last 20 years (articles published between 2005- 2025). Keywords used in the search included: BMI, Infertility, Insulin resistance, Obesity, PCOS.

Inclusion and Exclusion Criteria

This review included only peer-reviewed English articles reporting mean BMI, mean age, and PCOS prevalence. Studies were excluded if they lacked BMI or prevalence data, were non-research articles, had small sample sizes, or used unclear diagnostic criteria.

Study Selection and Data Extraction

The systematic review identified a total of 100 records through searches conducted in Google Scholar, Springer, and PubMed. After removing 15 duplicate studies, 85 records remained for screening. Based on title and abstract evaluation, 50 studies were selected for full-text assessment. Following the application of inclusion and exclusion criteria, 35 studies were reviewed in detail, and ultimately, 15 studies met the criteria for inclusion in this systematic review.




Figure 3: PRISMA flow diagram.

Results

The findings indicate a significant association between body mass index (BMI) and the prevalence of polycystic ovary syndrome (PCOS). Women with higher BMI values showed a greater likelihood of developing PCOS, with a particularly strong link observed in obese individuals. According to Alvarez-Blasco et al. (2006), the prevalence of PCOS in obese individuals was markedly higher compared to those with normal BMI, accompanied by increased insulin resistance and metabolic complications. Similarly, March et al. (2010) found that obesity significantly influenced symptom severity in PCOS patients, with a higher prevalence noted among individuals diagnosed using different criteria.

A global pattern of BMI-related PCOS prevalence was evident in studies conducted in different regions, including the United States, Spain, China, Iran, Australia, and Pakistan. For instance, Joshi et al. (2014) reported that younger women with higher BMI had an increased risk of PCOS, with obesity exacerbating hormonal imbalances and insulin resistance. Additionally, a study by Haase et al. (2023) highlighted that weight loss interventions in overweight and obese women with PCOS improved reproductive outcomes and reduced metabolic complications.

The review also indicated inconsistencies in PCOS prevalence across different studies, largely due to variations in diagnostic criteria. The Rotterdam criteria, National Institutes of Health (NIH) guidelines, and the National Institute of Child Health and Human Development (NICHD) criteria led to differences in reported prevalence rates. For example, Tehrani et al. (2011) observed a strong correlation between obesity and PCOS when using the Rotterdam criteria, while studies

Table 1: Summary of Key Studies on PCOS Prevalence and BMI.

PCOS= Polycystic ovarian syndrome, BMI= Body mass index, NICHD= National Institute of Child Health and Human Development, RC= Rotterdam

Author	Year	Country	Sample size of study	Mean age (yr)	Study Design	PCOS Diagnostic Criteria	Average BMI (kg/m ²)	Prevalence of PCOS (%)	Key Findings
Maria Trent et al. [32]	2005	USA	283	16.9	Cross-sectional study	NIH	31.7	34.3	Overweight girls with PCOS had a lower quality of life, and BMI played a significant role in mediating PCOS symptoms.
Alvarez-Blasco et al. [33]	2006	Spain	113	26	Observational study	NICHD	34.8	6.5	PCOS prevalence was significantly higher in obese individuals, with increased insulin resistance and metabolic complications.
X. Chen et al. [34]	2008	South China	915	31.8	Cross-sectional study	RC	20.94	2.4	PCOS prevalence was low in the population, with BMI levels significantly influencing hormonal and metabolic parameters.
Amini et al. [35]	2008	Iran	157	34.8	Case-control study	NIH	32.8	8.3	PCOS was more frequent in overweight/obese women with diabetes, with higher androgen levels and menstrual irregularities.
March et al. [36]	2010	South Australia	728	30.2	Population-based cohort study	NIH	25.7	17.8	PCOS prevalence varied depending on diagnostic criteria, with obesity playing a key role in symptom severity.
Tehrani et al. [37]	2011	Iran	1126	34.4	Community-based cross-sectional study	RC	26.2	14.6	Obesity was associated with increased PCOS prevalence, along with insulin resistance and dyslipidemia.
Hana Fakhoury et al. [38]	2012	Saudia Arabia	102	35.9	Cross-sectional study	RC	31.9	10	PCOS women showed significant hormonal imbalances, with BMI positively correlating with androgen levels.
Teede et al. [39]	2013	Australia	8612	30.5	Observational longitudinal study	Self-reported PCOS diagnosis	27.8	5.8	PCOS was linked to long-term weight gain, with metabolic syndrome risk increasing with BMI.
Joshi et al. [40]	2014	India	778	18.15	Cross-sectional study	RC	21.1	22.5	PCOS was prevalent in younger women, with obesity worsening symptoms and increasing insulin resistance.
Raziye Keskin Kurt et al. [41]	2014	Turkey	120	32.1	BMI-matched case-control study	RC	31.9	8.5	PCOS patients had higher inflammatory markers than BMI-matched controls, suggesting inflammation as a contributing factor.
Sharif et al. [42]	2016	Qatar	120	21	Cross-sectional study	NIH	22.95	11.7	PCOS prevalence was high among women with BMI >25kg/m ² , with increased risk of infertility.
E. Koivuaho et al. [43]	2019	Finland	280	46	Longitudinal cohort study	Self-reported PCOS diagnosis	25.75	4.45	Women with PCOS had increased long-term metabolic risks, with BMI strongly influencing outcomes.
Memon TF et al. [44]	2020	Pakistan	185	22.6	University-based cross-sectional study	RC	21.6	15.4	PCOS was highly prevalent among university students with BMI >25kg/m ² , with significant hormonal imbalances.
Neubronner et al. [45]	2021	Singapore	389	29.84	Prospective cross-sectional study	RC	25.14	34.4	PCOS phenotypes varied with BMI, with metabolic complications more common in obese individuals.
Haase et al. [46]	2023	UK	9955	27	Retrospective cohort study	EHR	31.2	10.5	Weight loss improved pregnancy chances in overweight PCOS women, with obesity linked to worse reproductive outcomes.

Discussion

The findings of this systematic review emphasize the critical role of BMI in the development and progression of PCOS. The evidence gathered from multiple studies confirms that women with higher BMI values are at a greater risk of developing PCOS, with metabolic disturbances and hormonal imbalances being more pronounced in overweight and obese individuals. Previous studies have also demonstrated that obesity contributes to hyperinsulinemia, which ex-

using NIH criteria demonstrated lower prevalence rates.

Overall, the results of this review confirm that PCOS is significantly influenced by BMI, with obesity being a key risk factor. The findings suggest that early interventions, particularly lifestyle modifications and weight management, could help in reducing the severity and prevalence of PCOS symptoms.

erbates androgen production and disrupts ovarian function, leading to PCOS symptoms (Tehrani et al., 2011; Fakhoury et al., 2012).

A key aspect highlighted in this review is the bidirectional relationship between PCOS and obesity. While obesity is recognized as a risk factor for PCOS, the syndrome itself can contribute to weight gain through mechanisms such as insulin resistance and metabolic dysfunction. This cycle perpetuates the severity of PCOS symptoms and

complicates management strategies (Haase et al., 2023). These findings underscore the need for targeted weight management strategies as a core component of PCOS treatment.

Another crucial factor discussed in this review is the variation in PCOS prevalence based on different diagnostic criteria. March et al. (2010) found that the application of the Rotterdam criteria resulted in higher prevalence rates compared to the NIH criteria, suggesting that broader diagnostic parameters may capture a wider spectrum of PCOS phenotypes. This inconsistency in diagnostic methods highlights the need for a standardized approach to improve comparability between studies.

Furthermore, the review highlights the impact of PCOS on metabolic health, with several studies reporting a higher incidence of insulin resistance, dyslipidemia, and metabolic syndrome in women with higher BMI. Studies such as those by Alvarez-Blasco et al. (2006) and Neubronner et al. (2021) indicated that obesity exacerbates metabolic complications in PCOS, increasing the risk of long-term health issues such as type 2 diabetes and cardiovascular diseases.

Additionally, regional differences in PCOS prevalence suggest that genetic, environmental, and lifestyle factors may influence the condition's manifestation. Studies from South Asia, such as those by Memon et al. (2020), reported a high prevalence of PCOS among university students, likely due to dietary patterns, sedentary lifestyles, and genetic predispositions. Conversely, studies conducted in Europe and North America, such as Haase et al. (2023), focused on the role of weight loss interventions in improving reproductive health among PCOS patients [32-46].

Conclusion

The systematic review confirms that BMI is a critical factor influencing PCOS prevalence. Overweight and obese women are at a higher risk of developing PCOS and experiencing more severe symptoms, including insulin resistance, hyperandrogenism, and metabolic syndrome. Given the heterogeneity in diagnostic criteria and study populations, future research should focus on developing standardized diagnostic methods and exploring targeted interventions for PCOS management.




Figure 4: Showing treatment measures for PCOS [31].

Limitations and Recommendations

This review is limited by restricted study access, variability in diagnostic criteria, and reliance on cross-sectional studies, which may introduce biases. Future research should prioritize longitudinal studies, standardized diagnostic criteria, and exploration of genetic, dietary, and lifestyle factors, as well as intervention-based studies on weight management and pharmacological treatments.

References

- Goodman NF, Cobin RH, Futterweit W, Glueck JS, Legro RS, et al. (2015) American Association of Clinical Endocrinologists, American College of Endocrinology, and androgen excess and PCOS society disease state clinical review: guide to the best practices in the evaluation and treatment of polycystic ovary syndrome-part 1. *Endocrine Practice* 21(11): 1291-1300.
- Okoroh EM, Hooper WC, Atrash HK, Yusuf HR, Boulet SL (2012) Prevalence of polycystic ovary syndrome among the privately insured, United States, 2003-2008. *American journal of obstetrics and gynecology* 207(4): 299-e1.
- Azziz R, Carmina E, Dewailly D, Diamanti-Kandarakis E, Escobar-Morreale HF, et al. (2009) The Androgen Excess and PCOS Society criteria for the polycystic ovary syndrome: the complete task force report. *Fertility and sterility* 91(2): 456-488.
- Legro RS (2000) The genetics of obesity Lessons for polycystic ovary syndrome. *Annals of the New York Academy of Sciences* 900(1): 193-202.
- Balen AH, Conway GS, Kaltsas G, Techatrasak K, Manning PJ, et al. (1995) Andrology: Polycystic ovary syndrome: the spectrum of the disorder in 1741 patients. *Human reproduction* 10(8): 2107-2111.
- Broekmans FJ, Knauff EA, Valkenburg O, Laven JS, Eijkemans MJ, et al. (2006) PCOS according to the Rotterdam consensus criteria: change in prevalence among WHO-II anovulation and association with metabolic factors. *BJOG: An International Journal of Obstetrics & Gynaecology* 113(10): 1210-1217.
- World Health Organization. Surveillance of chronic disease risk factors: country level data and comparable estimates.
- Carmina E (2004) Diagnosis of polycystic ovary syndrome: from NIH criteria to ESHRE-ASRM guidelines. *Minerva ginecologica* 56(1): 1-6.
- Azziz R, Carmina E, Chen Z, Dunaif A, Laven JS, et al. (2016) Polycystic ovary syndrome. *Nature reviews Disease primers* 2(1): 1-8.
- Diamanti-Kandarakis E, Dunaif A (2012) Insulin resistance and the polycystic ovary syndrome revisited: an update on mechanisms and implications. *Endocrine reviews* 33(6):981-1030.
- Alvarez-Blasco F, Botella-Carretero JJ, San Millán JL, Escobar-Morreale HF (2006) Prevalence and characteristics of the polycystic ovary syndrome in overweight and obese women. *Archives of internal medicine* 166(19): 2081-2086.
- Barber TM, Franks S (2021) Obesity and polycystic ovary syndrome. *Clinical endocrinology* 95(4): 531-541.
- Rich-Edwards JW, Spiegelman D, Garland M, Hertzmark E, Hunter DJ, et al. (2002) Physical activity, body mass index, and ovulatory disorder infertility. *Epidemiology* 13(2): 184-190.
- Beydoun HA, Stadtmauer L, Beydoun MA, Russell H, Zhao Y, et al. (2009) Polycystic ovary syndrome, body mass index and outcomes of assisted reproductive technologies. *Reproductive biomedicine online* 18(6): 856-863.
- Vague J (1956) The degree of masculine differentiation of obesities: a factor determining predisposition to diabetes, atherosclerosis, gout, and uric calculous disease. *The American journal of clinical nutrition* 4(1): 20-34.
- Lim SS, Davies MJ, Norman RJ, Moran LJ (2012) Overweight, obesity and central obesity in women with polycystic ovary syndrome: a systematic review and meta-analysis. *Human reproduction updates* 18(6): 618-637.
- DeBoer MD (2013) Obesity, systemic inflammation, and increased risk for cardiovascular disease and diabetes among adolescents: a need for screening tools to target interventions. *Nutrition* 29(2): 379-386.
- Willis DE, Franks ST (1995) Insulin action in human granulosa cells from normal and polycystic ovaries is mediated by the insulin receptor and not the type-I insulin-like growth factor receptor. *The Journal of Clinical Endocrinology & Metabolism* 80(12): 3788-3790.
- Chen X, Ni R, Mo Y, Li L, Yang D (2010) Appropriate BMI levels for PCOS patients in Southern China. *Human reproduction* 25(5): 1295-1302.
- Wild RA, Carmina E, Diamanti-Kandarakis E, Dokras A, Escobar-Morreale HF, et al. (2010) Assessment of cardiovascular risk and prevention of cardiovascular disease in women with the polycystic ovary syndrome: a consensus statement by the Androgen Excess and Polycystic Ovary Syndrome (AE-PCOS) Society. *The Journal of Clinical Endocrinology & Metabolism* 95(5): 2038-2049.
- Ehrmann DA, Liljenquist DR, Kasza K, Azziz R, Legro RS, et al. (2006) PCOS/Troglitazone Study Group. Prevalence and predictors of the metabolic syndrome in women with polycystic ovary syndrome. *The Journal of Clinical Endocrinology & Metabolism* 91(1): 48-53.
- Rosenfield RL (1999) Ovarian and adrenal function in polycystic ovary syndrome. *Endocrinology and metabolism clinics of North America* 28(2): 265-293.
- Littlejohn EE, Weiss RE, Deplewski D, Edidin DV, Rosenfield R (2007) Intractable early childhood obesity as the initial sign of insulin resistant hyperinsulinism and precursor of polycystic ovary syndrome. *Journal of Pediatric Endocrinology and Metabolism* 1: 41-52.
- Rosenfield RL (2007) Identifying children at risk for polycystic ovary syndrome. *The Journal of Clinical Endocrinology & Metabolism* 92(3): 787-796.
- Kaplowitz PB, Slora EJ, Wasserman RC, Pedlow SE, Herman-Giddens ME (2001) Earlier onset of puberty in girls: relation to increased body mass index and race. *Pediatrics* 108(2): 347-353.
- Lobo RA, Granger L, Goebelsmann UW, Manasco PK, Linsell C, et al. (1981) Elevations in unbound serum estradiol as a possible mechanism for inappropriate gonadotropin secretion in women with PCO. *The Journal of Clinical Endocrinology & Metabolism* 52(1): 156-158.
- Wehr E, Gruber HJ, Giuliani A, Möller R, Pieber TR, et al. (2011) The lipid accumulation product is associated with impaired glucose tolerance in PCOS women. *The Journal of Clinical Endocrinology & Metabolism* 96(6): E986-E990.
- Kahn HS (2005) The "lipid accumulation product" performs better than the body mass index for recognizing cardiovascular risk: a population-based comparison. *BMC cardiovascular disorders* 5(1): 1-0.
- Kahn HS (2006) The lipid accumulation product is better than BMI for identifying diabetes: a population-based comparison. *Diabetes care* 29(1): 151-153.
- Van Pelt RE, Evans EM, Schechtman KB, Ehsani AA, Kohrt WM (2002) Contributions of total and regional fat mass to risk for cardiovascular disease in older women. *American Journal of Physiology-Endocrinology and Metabolism*.
- Glueck CJ, Goldenberg N (2019) Characteristics of obesity in polycystic ovary syndrome: etiology, treatment, and genetics. *Metabolism* 92: 108-120.
- Trent M, Austin SB, Rich M, Gordon CM (2005) Overweight status of adolescent girls with polycystic ovary syndrome: body mass index as mediator of quality of life. *Ambul Pediatr* 5(2): 107-111.
- Alvarez-Blasco F, Botella-Carretero JJ, San Millán JL, Escobar-Morreale HF (2006) Prevalence and characteristics of the polycystic ovary syndrome in overweight and obese women. *Arch Intern Med* 166(19): 2081-2086.
- Chen X, Yang D, Mo Y, Li L, Chen Y, et al. (2008) Prevalence of polycystic ovary syndrome in unselected women from southern China. *Eur J Obstet Gynecol Reprod Biol* 139(1):59-64.
- Amini M, Horri N, Farmani M, Haghighi S, Sattari G, et al. (2008) Prevalence of polycystic ovary syndrome in reproductive-aged women with type 2 diabetes. *Gynecol Endocrinol* 24(8): 423-427.
- March WA, Moore VM, Willson KJ, Phillips DI, Norman RJ, et al. (2010) The prevalence of polycystic ovary syndrome in a community sample assessed under contrasting diagnostic criteria. *Hum Reprod* 25(2):544-551.
- Tehrani FR, Simbar M, Tohidi M, Hosseiniapanah F, Azizi F (2011)

- [The prevalence of polycystic ovary syndrome in a community sample of Iranian population: Iranian PCOS prevalence study. *Reprod Biol Endocrinol* 9: 39.](#)
38. [Fakhoury H, Tamim H, Ferwana M, Siddiqui IA, Adham M, et al. \(2012\) Age and BMI Adjusted Comparison of Reproductive Hormones in PCOS. *J Family Med Prim Care* 1\(2\):132-136.](#)
39. [Teede HJ, Joham AE, Paul E, Moran LJ, Loxton D, et al. \(2013\) Longitudinal weight gain in women identified with polycystic ovary syndrome: results of an observational study in young women. *Obesity \(Silver Spring\)* 21\(8\):1526-1532.](#)
40. [Joshi B, Mukherjee S, Patil A, Purandare A, Chauhan S, et al. \(2014\) A cross-sectional study of polycystic ovarian syndrome among adolescent and young girls in Mumbai, India. *Indian J Endocrinol Metab* 18\(3\): 317-324.](#)
41. [Keskin Kurt R, Okyay AG, Hakverdi AU \(2014\) The effect of obesity on inflammatory markers in patients with PCOS: a BMI-matched case-control study. *Arch Gynecol Obstet* 290: 315-319.](#)
42. [Sharif E, Rahman S, Zia Y, Rizk NM \(2016\) The frequency of polycystic ovary syndrome in young reproductive females in Qatar. *Int J Womens Health* 9:1-10.](#)
43. [Brower MA, Hai Y, Jones MR, Guo X, Chen YI, et al. \(2019\) Bidirectional Mendelian randomization to explore the causal relationships between body mass index and polycystic ovary syndrome. *Hum Reprod* 34\(1\):127-136.](#)
44. [Tariq Feroz Memon, Mehwish Channar, Syed Abdul Wadood Shah, Aiman Shaikh, Mehwish Batool et al. \(2020\) Polycystic Ovary Syndrome: Risk Factors and Associated Features among University Students in Pakistan. *Journal of Peoples University of Medical & Health Sciences Nawabshah. \(JPUMHS\)* 10\(1\).](#)
45. [Neubronner SA, Indran IR, Chan YH, Thu AWP, Yong EL \(2021\) Effect of body mass index \(BMI\) on phenotypic features of polycystic ovary syndrome \(PCOS\) in Singapore women: a prospective cross-sectional study. *BMC Womens Health* 21\(1\):135.](#)
46. [Haase CL, Varbo A, Laursen PN, Schnecke V, Balen AH \(2023\) Association between body mass index, weight loss and the chance of pregnancy in women with polycystic ovary syndrome and overweight or obesity: a retrospective cohort study in the UK. *Hum Reprod.* 2023 Mar 1;38\(3\):471-481.](#)